

APC/Fire™ 750 Rat IgG2b, κ Isotype Ctrl

Catalog # / Size:	2603350 / 100 µg 2603345 / 25 µg
Clone:	RTK4530
Isotype:	Rat IgG2b, κ
Immunogen:	Trinitrophenol + KLH
Preparation:	The antibody was purified by affinity chromatography and conjugated with APC/Fire&trade
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Workshop Number:	750 under optimal conditions.
Concentration:	0.2 mg/ml

Applications:

Applications: Flow Cytometry, Intracellular Staining for Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis as negative control. Use at concentrations comparable to those of the specific antibody of interest. * APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application Notes: The RTK4530 immunoglobulin is useful as an isotype-matched control (for the relevant formats) for Western blotting, immunoprecipitation, immunohistochemistry, functional assay, and immunofluorescence microscopy. The Ultra-LEAF™ purified antibody (Endotoxin < 0.01 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 400643, 400644, 400671, 400672, 400675, and 400676) as negative control.

Application References:

1. Cervantes-Barragan L, *et al.* 2007. *Blood* 109:1131.
2. Zeiser R, *et al.* 2007. *Blood* 109:2225.
3. Sasaki K, *et al.* 2008. *J. Immunol.* 181:104. [PubMed](#)
4. Duan J, *et al.* 2008. *P. Natl. Acad. Sci. USA* 105:5183. [PubMed](#)
5. Yi H, *et al.* 2009. *Blood* 113:5819. [PubMed](#)
6. Schafeer JS, *et al.* 2010. *J. Leukocyte Biol.* 87:301. [PubMed](#)
7. Lei GS, *et al.* 2015. *Infect Immun.* 83:572. [PubMed](#)
8. Richards J, *et al.* 2015. *Mol Cell Cardiol.* 79:21. [PubMed](#)

Description: The isotype of RTK4530 immunoglobulin is rat IgG2b, κ. This antibody was chosen as an isotype control after screening on a variety of resting, activated, live, and fixed mouse, rat and human tissues.